

## Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

**Web**: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Rabbit anti-Nogo Antibody

**Catalog:** DL93223A

**Synonyms:** KIAA0886; NOGO; Reticulon-4; Foocen; Neurite

outgrowth inhibitor; Nogo protein;

Neuroendocrine-specific protein; NSP;

Neuroendocrine-specific protein C homolog;

RTN-x: Reticulon-5

KLH-conjugated synthetic peptide encompassing a Immunogen:

sequence within the center region of human Nogo.

The exact sequence is proprietary.

Form: Liquid Concentration: 1mg/mL

**Size:** 100 u1/50 u1 Host: Rabbit



**Reactivity: Application:** WB, IHC

Human, Mouse, Rat, Bovine, Dog, M

onkey, Rabbit, Sheep

**Clonality:** Polyclonal **Dilution:** WB (1/500 - 1/1000),

IHC (1/100 - 1/200)

**Entrez Gene:** 57142/68585/83765

O9NOC3/O99P72/O9JK11 SwissProt:

**Purification:** The antibody was purified by immunogen affinity

chromatography.

**Buffer:** Liquid in 0.42% Potassium phosphate, 0.87%


Sodium chloride, pH 7.3, 30% glycerol, and 0.01%

sodium azide.

## **WB** description:

Western blot analysis of Nogo expression in HEK293T (A) whole cell lysates.





## **IHC** description:

Immunohistochemical analysis of Nogo staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

## **Storage:**

Store at -20°C. Avoid repeated freeze / thaw cycles.