

## Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

**Product name:** Rabbit anti-p65 Antibody

Catalog: DL93313A

**Synonyms:** HSN2; KDP; KIAA0344; PRKWNK1;

Serine/threonine-protein kinase WNK1; Erythrocyte

65 kDa protein; p65; Kinase deficient protein; Protein kinase lysine-deficient 1; Protein kinase

with no lysine 1; hWNK1

**Immunogen:** KLH-conjugated synthetic peptide encompassing a

sequence within the N-term region of human p65.

The exact sequence is proprietary.

Form: Liquid Concentration: 1mg/mL

Size: 100 ul/50 ul Host: Rabbit



**Reactivity: Application:** WB, IHC

Human, Mouse, Rat, Pig

**Clonality:** Polyclonal **Dilution:** WB (1/500 - 1/1000),

IHC (1/100 - 1/200)

**Entrez Gene:** 65125/232341/116477

O9H4A3/P83741/O9JIH7 SwissProt:

**Purification:** The antibody was purified by immunogen affinity

chromatography.

**Buffer:** Liquid in 0.42% Potassium phosphate, 0.87%


Sodium chloride, pH 7.3, 30% glycerol, and 0.01%

sodium azide

**WB** description:

Western blot analysis of p65 expression in HEK293T (A), HeLa (B) whole cell lysates.





## **IHC** description:

Immunohistochemical analysis of p65 staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

## **Storage:**

Store at -20°C. Avoid repeated freeze / thaw cycles.