Manufacturer/Supplier:
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District Wuxi Jiangsu Province, China
TELE：86-510-82732223 FAX：86-510-82720101-8014
Web : www.dldevelop.com
Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Rabbit anti-Caspase 6 Antibody
Catalog: DL93358A

Synonyms: MCH2; Caspase-6; CASP-6; Apoptotic protease Mch-2

Immunogen: KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Caspase 6. The exact sequence is proprietary.

Form: Liquid **Concentration:** 1mg/mL

Size: 100 ul/50 ul **Host:** Rabbit

Reactivity: Human,Mouse,Rat,Bovine,Monkey **Application:** WB, IHC, IF/IC

Clonality: Polyclonal

Dilution: WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500)

Entrez Gene: 839/12368

SwissProt: P55212/O08738/O35397

Purification: The antibody was purified by immunogen affinity chromatography.

Buffer: Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of Caspase 6 expression in HepG2 (A), Raw264.7 (B) whole cell lysates.



IHC description:

Immunohistochemical analysis of Caspase 6 staining in human prostate cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IF/ICC description:

Immunofluorescent analysis of Caspase 6 staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary anti



Storage:

Store at -20℃. Avoid repeated freeze / thaw cycles.