

Address: A1-203 I	SCI&TECH DEVELOPMENT CO., LTD Mingpin CityII,No.8 Xihudong Road.Liangxi District		
Wuxi Jiangsu Province, China TELE: 86-510-82732223 FAX: 86-510-82720101-8014			
Web : www.dldevelop.com FAX: 30-510-32720101-3014			
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>			
Product name:	Rabbit anti-PKR Antibody		
Catalog:	DL93395A		
Synonyms:	PKR; PRKR; Interferon-induced, double-stranded RNA-activated protein kinase; Eukaryotic translation initiation factor 2-alpha kinase 2; eIF-2A protein kinase 2; Interferon-inducible RNA-dependent protein kinase; P1/eIF-2A protein kinase; Protein kinase RN		
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human PKR. The exact sequence is proprietary.		
Form: Liquid	Concentration: 1mg/mL		
Size: 100 ul/50 ul	Host: Rabbit		




Reactivity: Human, Mouse, Rat		Application: WB, IHC
Clonality: Polyclonal		Dilution: WB (1/500 - 1/1000), IHC (1/100 - 1/200)
Entrez Gene:	5610/19106/54287	
SwissProt:	P19525/Q03963/Q63184	
Purification:	The antibody was purified by immunogen affinity chromatography.	
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.	

WB description:

Western blot analysis of PKR expression in THP1 (A), NS-1 (B), PC12 (C) whole cell lysates.

DEVELOP



IHC description:

Immunohistochemical analysis of PKR staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.