

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

Web : www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Rabbit anti-TRAP220 (pT1457) Antibody

Catalog: DL93460A

Synonyms: ARC205; CRSP1; CRSP200; DRIP205; DRIP230;
PBP; PPARBP; PPARGBP; RB18A; TRAP220;
TRIP2; Mediator of RNA polymerase II
transcription subunit 1; Activator-recruited cofactor
205 kDa component; ARC205; Mediator complex
subunit 1; Peroxisome proliferator-activ

Immunogen: KLH-conjugated synthetic peptide encompassing a
sequence within the C-term region of human
TRAP220 (pT1457). The exact sequence is
proprietary.

Form: Liquid

Concentration: 1mg/mL

Size: 100 ul/50 ul

Host: Rabbit

Reactivity:
Human, Mouse, Rat, Monkey

Application: WB, IHC, IF/IC, IP,
ChIP

Clonality: Polyclonal

Dilution: WB (1/500 - 1/1000),
IHC (1/100 - 1/200), IF/IC (1/100 -
1/500), IP (1/10 - 1/100), ChIP
(Use at an assay dependent
concentration)

Entrez Gene: 5469/19014


SwissProt: Q15648/Q925J9

Purification: The antibody was purified by immunogen affinity
chromatography.

Buffer: Liquid in 0.42% Potassium phosphate, 0.87%
Sodium chloride, pH 7.3, 30% glycerol, and 0.01%
sodium azide.

WB description:

Western blot analysis of TRAP220 (pT1457) expression in Jurkat
(A), HeLa (B), NIH3T3 (C) whole cell lysates.




IHC description:

Immunohistochemical analysis of TRAP220 (pT1457) staining in human colon cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IF/ICC description:

Immunofluorescent analysis of TRAP220 (pT1457) staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the prim



IP description:

ChIP description:

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.