

Manufacturer/Supplier:			
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD			
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District			
Wuxi Jiangsu Province, China			
TELE: 86-510-82732223 FAX: 86-510-82720101-8014			
Web : <u>www.dldevelop.com</u>			
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>			
Product name:	Rabbit anti-Alpha-synuclein (pY133) Antibody		
Catalog:	DL93524A		
U			
Synonyms:	NACP; PARK1; Alpha-synuclein; Non-A beta		
	component of AD amyloid; Non-A4 component of		
	amyloid precursor; NACP		
	uniyola piccusol, tatel		
Immunogen:	Immunogen: KLH-conjugated synthetic peptide encompassing a		
immunogen.			
	sequence within the C-term region of human Alpha-synuclein (pY133). The exact sequence is		
	proprietary.		
Form: Liquid	Concentration: 1mg/mL		
Size: 100 ul/50 ul	Host: Rabbit		
Reactivity:	Application: WB, IHC		

DEVELOP.


Human,Mouse,Rat,Bovine,Monke y,Pig,Rabbit,Sheep

Clonality: Polyclo	nal	Dilution: WB (1/500 - 1/1000), IHC (1/100 - 1/200)	
Entrez Gene:	6622/20617/292	19	
SwissProt:	P37840/O55042/P37377		
Purification:	The antibody was purified by immunogen affinity chromatography.		
Buffer:	-	Potassium phosphate, 0.87% , pH 7.3, 30% glycerol, and 0.01%	

WB description:

Western blot analysis of Alpha-synuclein (pY133) expression in HEK293T (A) whole cell lysates.

DEVELOP



IHC description:

Immunohistochemical analysis of Alpha-synuclein (pY133) staining in human prostate cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

