

Manufacturer/Supplier:			
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD			
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District			
Wuxi Jiangsu Province, China			
TELE: 86-510-82732223 FAX: 86-510-82720101-8014			
Web : <u>www.dldevelop.com</u>			
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>			
Product name:	Rabbit anti-Tyrosine Hydroxylase (pS19) Antibody		
Catalog:	DL93531A		
Synonyms:	TYH; Tyrosine 3-monooxygenase; Tyrosine		
	3-hydroxylase; TH		
Immunogen:	KLH-conjugated synthetic peptide encompassing a		
_	sequence within the N-term region of human		
	Tyrosine Hydroxylase (pS19). The exact sequence		
	is proprietary.		
Form: Liquid		Concentr	ation: 1mg/mL
1			C
Size: 100 ul/50 ul		Host: Rab	bit
<b>Reactivity:</b>		Applicati	on: WB, IHC, IF/IC
Human,Mouse,Rat,Bovine,Dog			




**Clonality:** Polyclonal

**Dilution:** WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500)


Entrez Gene:7054/21823/25085SwissProt:P07101/P24529/P04177Purification:The antibody was purified by immunogen affinity<br/>chromatography.Buffer:Liquid in 0.42% Potassium phosphate, 0.87%<br/>Sodium chloride, pH 7.3, 30% glycerol, and 0.01%<br/>sodium azide.

#### **WB description:**

Western blot analysis of Tyrosine Hydroxylase (pS19) expression in HEK293T (A), A549 (B) whole cell lysates.



# **DEVELOP**




#### **IHC description:**

Immunohistochemical analysis of Tyrosine Hydroxylase (pS19) staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

#### **IF/ICC description:**

Immunofluorescent analysis of Tyrosine Hydroxylase (pS19) staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with

## **DEVELOP**.



### Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

