

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82732223		FAX:	86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name:	Rabbit anti-WASP (pY290) Antibody			
Catalog:	DL93544A			
8				
Synonyms:	IMD2; Wiskott-Aldrich syndrome protein; WASp			
Synonyms.	1101D2, 11 15K0tt	r Harlen Syr	laronie protein, wriop	
Immunogen:	KLH-conjugated synthetic peptide encompassing a			
ininunogen.	sequence within the center region of human WASP			
	(pY290). The exact sequence is proprietary.			
T. T. 1		C ((* 1 / T	
Form: Liquid		Concentr	ation: 1mg/mL	
C* 100 1/50 1			1.1	
Size: 100 ul/50 ul		Host: Rat	obit	
Reactivity: Human, Mouse, Rat		Applicati	on: WB, IHC, IF/IC, FC	
Classifier Debuteral Dilations WD (1/500 - 1/1000)			WD(1/500-1/1000)	
Clonality: Polyclonal		Dilution: WB (1/500 - 1/1000),		
		IHC (1/10	0 - 1/200), IF/IC (1/100 -	




1/500), FC (1/100 - 1/200)

Entrez Gene:	7454/22376
SwissProt:	P42768/P70315
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of WASP (pY290) expression in Jurkat (A), TF1 (B) whole cell lysates.




IHC description:



Immunohistochemical analysis of WASP (pY290) staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IF/ICC description:

Immunofluorescent analysis of WASP (pY290) staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary a



FC description:



Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

