

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Rabbit anti-CD88 Antibody

Catalog: DL93586A

Synonyms: C5AR; C5R1; C5a anaphylatoxin chemotactic

receptor 1; C5a anaphylatoxin chemotactic receptor;

C5a-R; C5aR; CD88

Immunogen: KLH-conjugated synthetic peptide encompassing a

sequence within the C-term region of human CD88.

The exact sequence is proprietary.

Form: Liquid Concentration: 1mg/mL

Size: 100 u1/50 u1 Host: Rabbit

Reactivity: Human, Mouse, Rat **Application:** WB, IHC, IF/IC



Clonality: Polyclonal **Dilution:** WB (1/500 - 1/1000),

IHC (1/100 - 1/200), IF/IC (1/100 -

1/500)

Entrez Gene: 728/113959

P21730/P30993/P97520 SwissProt:

Purification: The antibody was purified by immunogen affinity

chromatography.

Buffer: Liquid in 0.42% Potassium phosphate, 0.87%


Sodium chloride, pH 7.3, 30% glycerol, and 0.01%

sodium azide.

WB description:

Western blot analysis of CD88 expression in HepG2 (A), mouse brain (B), rat heart (C) whole cell lysates.






IHC description:

Immunohistochemical analysis of CD88 staining in human kidney formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IF/ICC description:

Immunofluorescent analysis of CD88 staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody





Storage: Store at -20°C. Avoid repeated freeze / thaw cycles.