

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82732223 FAX: 86-510-82720101-8014				
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name: Catalog:	Rabbit anti-Cy DL93708A	vtokeratin 8 (J	oS432) Antibody	
Synonyms:	CYK8; Keratin Cytokeratin-8; Kb8		oskeletal 8; in-8; K8; Type-II keratin	
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Cytokeratin 8. The exact sequence is proprietary.			
Form: Liquid		Concentra	ation: 1mg/mL	
Size: 100 ul/50 ul		Host: Rab	bit	
Reactivity: Human, Mouse		Applicatio	Application: WB, IHC, IF/IC	




**Clonality:** Polyclonal

**Dilution:** WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500)


Entrez Gene:	3856/16691
SwissProt:	P05787/P11679
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

#### **WB description:**

Western blot analysis of Cytokeratin 8 (pS432) expression in H522 (A), H1792 (B), mouse pancreatic cancer (C) whole cell lysates.



# **DEVELOP**




#### **IHC description:**

Immunohistochemical analysis of Cytokeratin 8 (pS432) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

#### **IF/ICC description:**

Immunofluorescent analysis of Cytokeratin 8 (pS432) staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the p

## **DEVELOP**.



### Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

