

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

Web : www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Rabbit anti-Ezrin/Radixin/Moesin (pT567/564/558)
Antibody

Catalog: DL93742A

Synonyms: MSN; Moesin; Membrane-organizing extension
spike protein; RDX; Radixin; EZR; VIL2; Ezrin;
Cytovillin; Villin-2; p81

Immunogen: KLH-conjugated synthetic peptide encompassing a
sequence within the C-term region of human
Ezrin/Radixin/Moesin . The exact sequence is
proprietary.

Form: Liquid

Concentration: 1mg/mL

Size: 100 ul/50 ul

Host: Rabbit

Reactivity:

Human, Mouse, Rat, Bovine, Monkey, Pig, Zebrafish

Application: WB, IHC, IF/IC**Clonality:** Polyclonal

Dilution: WB (1/500 - 1/1000),
IHC (1/100 - 1/200), IF/IC (1/100 - 1/500)

Entrez Gene:

17698; 19684; 22350/81521; 54319

SwissProt:

P26038; P35241; P15311/P26041; P26043;
P26040/O35763; P31977

Purification:


The antibody was purified by immunogen affinity chromatography.

Buffer:

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of Ezrin/Radixin/Moesin (pT567/564/558) expression in SKOV3 (A), NIH3T3 (B) whole cell lysates.




IHC description:

Immunohistochemical analysis of Ezrin/Radixin/Moesin (pT567/564/558) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IF/ICC description:

Immunofluorescent analysis of Ezrin/Radixin/Moesin (pT567/564/558) staining in SKOV3 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were p



Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.