

Manufaaturar/Su	nnlior			
Manufacturer/Supplier: WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82	2732223	FAX:	86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name:	Rabbit anti-P2Y4 Antibody			
Catalog:	DL93763A			
cg.	22,0,0011			
Synonyms:	NDLL D2V numino conton 4, D2V4, D2D, Uniding			
Synonyms.	NRU; P2Y purinoceptor 4; P2Y4; P2P; Uridine			
	nucleotide recep	tor; UNK		
-				
Immunogen:	KLH-conjugated synthetic peptide encompassing a			
	sequence within the center region of human P2Y4.			
	The exact sequence is proprietary.			
Form: Liquid		Concentr	ation: 1mg/mL	
Size: 100 ul/50 ul		Host: Rabbit		
Reactivity: Human, Mouse, Rat		Annlicati	on: WB, IF/IC	
iteacer ny ritaina	ii,iii us 0,1 (ut	ppneut		
Clonality: Polyclonal		Dilution: WB (1/500 - 1/1000),		
Cionanty, i orycional Dilution, wb			(1/300 - 1/1000),	

www.dldevelop.com




IF/IC (1/100 - 1/500)

Entrez Gene:	5030/57385/63843
SwissProt:	P51582/Q9JJS7/O35811
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:


Western blot analysis of P2Y4 expression in HeLa (A), rat brain (B), rat lung (C) whole cell lysates.



IF/ICC description:



Immunofluorescent analysis of P2Y4 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody i



Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

