

## Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

**Product name:** Rabbit anti-ZNF265 Antibody

Catalog: DL93964A

**Synonyms:** ZIS; ZNF265; Zinc finger Ran-binding

domain-containing protein 2; Zinc finger protein

265; Zinc finger, splicing

Immunogen: KLH-conjugated synthetic peptide encompassing a

sequence within the center region of human ZNF265. The exact sequence is proprietary.

Form: Liquid Concentration: 1mg/mL

Size: 100 ul/50 ul Host: Rabbit

Reactivity: Application: WB, IHC, IF/IC

Human, Mouse, Rat, Chicken, Dog,



# Monkey, Pig, Rabbit

**Clonality:** Polyclonal **Dilution:** WB (1/500 - 1/1000),

IHC (1/100 - 1/200), IF/IC (1/100 -

1/500)

**Entrez Gene:** 9406/53861/58821

O95218/O9R020/O35986 SwissProt:

**Purification:** The antibody was purified by immunogen affinity

chromatography.

**Buffer:** Liquid in 0.42% Potassium phosphate, 0.87%


Sodium chloride, pH 7.3, 30% glycerol, and 0.01%

sodium azide

#### **WB** description:

Western blot analysis of ZNF265 expression in HepG2 (A), MCF7 (B) whole cell lysates.






### **IHC** description:

Immunohistochemical analysis of ZNF265 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

## **IF/ICC description:**

Immunofluorescent analysis of ZNF265 staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibod





**Storage:** 

Store at -20°C. Avoid repeated freeze / thaw cycles.