

## Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>

**Product name:** Rabbit anti-MAPKAPK3 Antibody

Catalog: DL93978A

**Synonyms:** MAP kinase-activated protein kinase 3;

MAPK-activated protein kinase 3; MAPKAP kinase

3; MAPKAP-K3; MAPKAPK-3; MK-3;

Chromosome 3p kinase; 3pK

Immunogen: KLH-conjugated synthetic peptide encompassing a

sequence within the C-term region of human

MAPKAPK3. The exact sequence is proprietary.

Form: Liquid Concentration: 1mg/mL

Size: 100 ul/50 ul Host: Rabbit

**Reactivity:** Human,Pig Application: WB, IHC, IF/IC



**Clonality:** Polyclonal **Dilution:** WB (1/500 - 1/1000),

IHC (1/100 - 1/200), IF/IC (1/100 -

1/500)

**Entrez Gene:** 7867

Q16644 SwissProt:

**Purification:** The antibody was purified by immunogen affinity

chromatography.

**Buffer:** Liquid in 0.42% Potassium phosphate, 0.87%


Sodium chloride, pH 7.3, 30% glycerol, and 0.01%

sodium azide.

## **WB** description:

Western blot analysis of MAPKAPK3 expression in Hela (A), Jurkat (B) whole cell lysates.






## **IHC** description:

Immunohistochemical analysis of MAPKAPK3 staining in human heart formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

## **IF/ICC description:**

Immunofluorescent analysis of MAPKAPK3 staining in Hela cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibo





**Storage:** Store at -20°C. Avoid repeated freeze / thaw cycles.