

Manufacturer/Supplier:	
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD	
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District	
Wuxi Jiangsu Province, China	
TELE: 86-510-82	FAX: 86-510-82720101-8014
Web : <u>www.dldevelop.com</u>	
Email: info@dldevelop.com.cn service@dldevelop.com.cn	
Product name:	Rabbit anti-CstF-64T Antibody
Catalog:	DL94004A
Synonyms:	KIAA0689; Cleavage stimulation factor subunit 2
	tau variant; CF-1 64 kDa subunit tau variant;
	Cleavage stimulation factor 64 kDa subunit tau
	variant; CSTF 64 kDa subunit tau variant;
	TauCstF-64
Immunogen:	KLH-conjugated synthetic peptide encompassing a
	sequence within the center region of human
	CstF-64T. The exact sequence is proprietary.
Form: Liquid	Concentration: 1mg/mL
Form. Elquid	Concentration. Ting/IIL
Size: 100 ul/50 ul	Host: Rabbit



Reactivity: Human.Mouse.Rat.Bovine

Application: WB, IHC


Clonality: Polyclonal Dilution: WB (1/500 - 1/1000), IHC (1/100 - 1/200) **Entrez Gene:** 23283/83410 O9H0L4/O8C7E9 SwissProt: **Purification:** The antibody was purified by immunogen affinity chromatography. **Buffer:** Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide

WB description:

Western blot analysis of CstF-64T expression in Hela (A), PC3 (B), MCF7 (C) whole cell lysates.



DEVELOP



IHC description:

Immunohistochemical analysis of CstF-64T staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.