

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

Web : www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Rabbit anti-p47 phox (pS345) Antibody

Catalog: DL94058A

Synonyms: NOXO2; SH3PXD1A; Neutrophil cytosol factor 1;
NCF-1; 47 kDa autosomal chronic granulomatous
disease protein; 47 kDa neutrophil oxidase factor;
NCF-47K; Neutrophil NADPH oxidase factor 1;
Nox organizer 2; Nox-organizing protein 2; SH3
and PX domain-contain

Immunogen: KLH-conjugated synthetic peptide encompassing a
sequence within the C-term region of human p47
phox. The exact sequence is proprietary.

Form: Liquid

Concentration: 1mg/mL

Size: 100 ul/50 ul

Host: Rabbit

Reactivity: Human, Mouse

Application: WB, IHC

Clonality: Polyclonal

Dilution: WB (1/500 - 1/1000),
IHC (1/100 - 1/200)

Entrez Gene: 653361/17969


SwissProt: P14598/Q09014

Purification: The antibody was purified by immunogen affinity chromatography.

Buffer: Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of p47 phox (pS345) expression in HepG2 (A), NIH3T3 (B) whole cell lysates.



IHC description:

Immunohistochemical analysis of p47 phox (pS345) staining in human lymph node formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.