

Manufacturer/Supplier:			
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD			
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District			
Wuxi Jiangsu Province, China			
TELE: 86-510-82732223 FAX: 86-510-82720101-8014			
Web : <u>www.dldevelop.com</u>			
Email: info@dldevelop.com.cn service@dldevelop.com.cn			
Product name:	Rabbit anti-JAK2 Antibody		
Catalog:	DL94112A		
Synonyms:	Tyrosine-protein kinase JAK2; Janus kinase 2;		
	JAK-2		
Immunogen:	KLH-conjugated synthetic peptide encompassing a		
8	sequence within the center region of human JAK2.		
	The exact sequence is proprietary.		
Form: Liquid		Concentra	ation: 1mg/mL
Size: 100 ul/50 ul		Host: Rab	bit
Desetivity		Amplicati	
Reactivity:	Dec Mentere	Application	on: WB, IHC, IF/IC
Human,Mouse,Rat,Dog,Monkey,			
Pig,Rabbit			



Clonality: Polyclonal

Dilution: WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500)


- **Entrez Gene:** 3717/16452/24514
- SwissProt: 060674/Q62120/Q62689
- **Purification:** The antibody was purified by immunogen affinity chromatography.
- **Buffer:** Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of JAK2 expression in Jurkat (A), K562 (B), HEK293T (C) whole cell lysates.



DEVELOP




IHC description:

Immunohistochemical analysis of JAK2 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IF/ICC description:

Immunofluorescent analysis of JAK2 staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody

DEVELOP.



Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

