

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82732223		FAX:	86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: info@dldevelop.com.cn service@dldevelop.com.cn				
Product name:	Rabbit anti-Der	matin Antibo	ody	
Catalog:	DL94116A			
Synonyms:	DMT; EPB49; Dematin; Dematin actin-binding protein; Erythrocyte membrane protein band 4.9			
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human			
	Dematin. The e	exact sequen	ce is proprietary.	
Form: Liquid		Concentr	ation: 1mg/mL	
Size: 100 ul/50 ul		Host: Rab	bbit	
<b>Reactivity:</b> Human,Mouse,Boy	vine	Applicati	on: WB, IHC, IF/IC	




**Clonality:** Polyclonal

**Dilution:** WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500)


Entrez Gene:	2039/13829
SwissProt:	Q08495/Q9WV69
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

#### **WB description:**

Western blot analysis of Dematin expression in Jurkat (A), HEK293T (B) whole cell lysates.



# **DEVELOP**




## **IHC description:**

Immunohistochemical analysis of Dematin staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

#### **IF/ICC description:**

Immunofluorescent analysis of Dematin staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antib

# **DEVELOP**.



## Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

