

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82	2732223	FAX:	86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name:	Rabbit anti-DYRK1B Antibody			
Catalog:	DL94164A			
Synonyms:	MIRK; Dual specificity			
	tyrosine-phosp	phorylation-re	egulated kinase 1B;	
	Minibrain-rela	ated kinase; N	firk protein kinase	
Immunogen:	KLH-conjugated synthetic peptide encompassing a			
	sequence within the center region of human			
	DYRK1B. The exact sequence is proprietary.			
Form: Liquid		Concentra	ation: 1mg/mL	
C* 100 1/50 1			1.1.	
Size: 100 ul/50 ul		Host: Rab	obit	
Depativity		Applicati	WD IUC	
Reactivity:Application: WB, IHC			UII. WD , IIIC	
Human,Mouse,Bovine				




Clonality: Polyclonal

Dilution: WB (1/500 - 1/1000), IHC (1/100 - 1/200)


Entrez Gene:	9149/13549
SwissProt:	Q9Y463/Q9Z188
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of DYRK1B expression in Hela (A), HT1080 (B) whole cell lysates.



DEVELOP



IHC description:

Immunohistochemical analysis of DYRK1B staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.