

## Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

**Web**: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Rabbit anti-Beta-1 Adrenergic Receptor Antibody

**Catalog:** DL94170A

**Synonyms:** ADRB1R; B1AR; Beta-1 adrenergic receptor;

Beta-1 adrenoreceptor; Beta-1 adrenoceptor

Immunogen: KLH-conjugated synthetic peptide encompassing a

sequence within the center region of human Beta-1

Adrenergic Receptor. The exact sequence is

proprietary.

Form: Liquid Concentration: 1mg/mL

**Size:** 100 ul/50 ul Host: Rabbit

**Reactivity:** Human **Application:** WB, IHC, IF/IC



**Clonality:** Polyclonal **Dilution:** WB (1/500 - 1/1000),

IHC (1/100 - 1/200), IF/IC (1/100 -

1/500)

Entrez Gene: 153

P08588 SwissProt:

**Purification:** The antibody was purified by immunogen affinity

chromatography.

**Buffer:** Liquid in 0.42% Potassium phosphate, 0.87%


Sodium chloride, pH 7.3, 30% glycerol, and 0.01%

sodium azide.

**WB** description:

Western blot analysis of Beta-1 Adrenergic Receptor expression in HepG2 (A), A431 (B) whole cell lysates.






## **IHC** description:

Immunohistochemical analysis of Beta-1 Adrenergic Receptor staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

## **IF/ICC description:**

Immunofluorescent analysis of Beta-1 Adrenergic Receptor staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with





**Storage:** Store at -20°C. Avoid repeated freeze / thaw cycles.