

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82	732223	FAX:	86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name:	Rabbit anti-NDRG4 Antibody			
Catalog:	DL94226A			
Synonyms:	BDM1; KIAA1180; Protein NDRG4; Brain			
	development-re	elated molec	ule 1; N-myc	
	downstream-re	gulated gene	4 protein; Vascular	
	smooth muscle	cell-associa	ted protein 8; SMAP-8	
Immunogen:	KLH-conjugated synthetic peptide encompassing a			
	sequence withi	n the center	region of human	
	NDRG4. The e	exact sequence	e is proprietary.	
Form: Liquid		Concentra	ation: 1mg/mL	
Size: 100 ul/50 ul		Host: Rab	bit	
Reactivity: Human, Mouse, Rat		Application	Application: WB, IHC	




**Clonality:** Polyclonal

**Dilution:** WB (1/500 - 1/1000), IHC (1/100 - 1/200)


Entrez Gene:	65009/234593
SwissProt:	Q9ULP0/Q8BTG7/Q9Z2L9
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

## **WB description:**

Western blot analysis of NDRG4 expression in Jurkat (A), HepG2 (B) whole cell lysates.



## **DEVELOP**



## **IHC description:**

Immunohistochemical analysis of NDRG4 staining in human liver formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. AEC was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

## **Storage:**

Store at -20°C. Avoid repeated freeze / thaw cycles.