

Manufacturer/Su				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82732223 FAX: 86-510-82720101-8014				
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name:	Rabbit anti-Ubiquilin 4 Antibody			
Catalog:	DL94244A			
C				
Synonyms:	Clorf6; CIP75; UBIN; Ubiquilin-4; Ataxin-1			
			otein; A1Up; Ataxin-1	
	ubiquitin-like-ir	· ·	· · ·	
	1	01	tein of 75 kDa; CIP75	
Immunogen:	KLH-conjugate	d synthetic	peptide encompassing a	
g			region of human	
	1		ence is proprietary.	
	Obiquinii 4. Th	e exact sequ	tenee is proprietary.	
Form: Liquid		Concontr	ation: 1mg/mL	
Form. Elquid		Concentra	ation. mig/mL	
Size: 100 ul/50 ul		Host: Rab	h;t	
Size: 100 ul/30 ul		nost: Kau	oon	
Reactivity: Human, Mouse Application: WB, IHC			WD HIC	
Reactivity: Human, Mouse		Application	on: wB, IHC	



Clonality: Polyclonal

Dilution: WB (1/500 - 1/1000), IHC (1/100 - 1/200)


Entrez Gene:	56893/94232
SwissProt:	Q9NRR5/Q99NB8
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of Ubiquilin 4 expression in COS7 (A), HEK293T (B), AML12 (C), PC12 (D) whole cell lysates.



DEVELOP



IHC description:

Immunohistochemical analysis of Ubiquilin 4 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.