

Manufacturer/Su	nnlier·				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD					
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District					
Wuxi Jiangsu Province, China					
TELE: 86-510-82732223 FAX: 86-510-82720101-8014					
Web : www.dldevelop.com					
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>					
Product name: Catalog:	Rabbit anti-Alp DL94378A	ha-synucleii	n (pY125) Antibody		
Synonyms:	NACP; PARK1; Alpha-synuclein; Non-A beta component of AD amyloid; Non-A4 component of amyloid precursor; NACP				
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Alpha-synuclein. The exact sequence is proprietary.				
Form: Liquid		Concentra	ation: 1mg/mL		
Size: 100 ul/50 ul		Host: Rab	bit		
Reactivity:Application: WB, IHCHuman,Mouse,Rat,Bovine,Monke					



y,Pig


Clonality: Polyclo		Dilution: WB (1/500 - 1/2000), IHC (1/50 - 1/200)	
Entrez Gene:	6622/20617/2921	9	
SwissProt:	P37840/O55042/P37377		
Purification:	The antibody was chromatography.	purified by immunogen affinity	
Buffer:	1	Potassium phosphate, 0.87% pH 7.3, 30% glycerol, and 0.01%	

WB description:

Western blot analysis of Alpha-synuclein (pY125) expression in HeLa (A), mouse brain (B), rat brain (C) whole cell lysates.



DEVELOP



IHC description:

Immunohistochemical analysis of Alpha-synuclein (pY125) staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

