

Manufacturer/Supplier:		
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD		
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District		
Wuxi Jiangsu Province, China		
TELE: 86-510-82732223 FAX: 86-510-82720101-8014		
Web : <u>www.dldevelop.com</u>		
Email: info@dldevelop.com.cn service@dldevelop.com.cn		
Product name: Catalog:	Rabbit anti-ZNF392 Antibody DL94473A	
Synonyms:	ZNF392; ZNF435; Zinc finger and SCAN domain-containing protein 16; Zinc finger protein 392; Zinc finger protein 435	
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human ZNF392. The exact sequence is proprietary.	
Form: Liquid	Concentration: 1mg/mL	
Size: 100 ul/50 ul	Host: Rabbit	
Reactivity: Human,Mouse,Rat	Application: WB, IHC, IF/IC	



Clonality: Polyclonal

Dilution: WB (1/500 - 1/1000), IHC (1/50 - 1/100), IF/IC (1/100 - 1/500)


Entrez Gene:	80345
SwissProt:	Q9H4T2
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of ZNF392 expression in HCT116 (A), DLD (B), mouse brain (C) whole cell lysates.



DEVELOP



IHC description:


Immunohistochemical analysis of ZNF392 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IF/ICC description:

Immunofluorescent analysis of ZNF392 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody



DEVELOP.



Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

