

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Rabbit anti-TBK1 Antibody

Catalog: DL94544A

NAK; Serine/threonine-protein kinase TBK1; **Synonyms:**

NF-kappa-B-activating kinase; T2K;

TANK-binding kinase 1

Immunogen: KLH-conjugated synthetic peptide encompassing a

sequence within the center region of human TBK1.

The exact sequence is proprietary.

Form: Liquid Concentration: 1mg/mL

Size: 100 u1/50 u1 Host: Rabbit

Reactivity: Human, Mouse, Rat **Application:** WB, IHC



Clonality: Polyclonal **Dilution:** WB (1/500 - 1/1000),

IHC (1/50 - 1/100)

Entrez Gene: 29110/56480

O9UHD2/O9WUN2 SwissProt:

Purification: The antibody was purified by immunogen affinity

chromatography.

Buffer: Liquid in 0.42% Potassium phosphate, 0.87%


Sodium chloride, pH 7.3, 30% glycerol, and 0.01%

sodium azide.

WB description:

Western blot analysis of TBK1 expression in mouse testis (A), rat testis (B), U87MG (C), A549 (D) whole cell lysates.





IHC description:

Immunohistochemical analysis of TBK1 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.