

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82	732223	FAX:	86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name:	Rabbit anti-Synaptotagmin (pT202/199) Antibody			
Catalog:	DL94736A			
Synonyms:	SYT1; SVP65; SYT; Synaptotagmin-1; Synaptotagmin I; SytI; p65; SYT2;			
	Synaptotagmin-	2; Synaptot	agmin II; SytII	
Immunogen:	KLH-conjugated synthetic peptide encompassing a			
	sequence within the center region of human			
	Synaptotagmin. The exact sequence is proprietary.			
Form: Liquid		Concentr	ation: 1mg/mL	
Size: 100 ul/50 ul		Host: Rab	obit	
Reactivity: Human, Mouse, Rat Application: WB, IHC				
Reactivity: Humai	n,mouse,Kat	Applicati	on: wB, IHC	



Clonality: Polyclonal

Dilution: WB (1/500 - 1/1000), IHC (1/50 - 1/100)


Entrez Gene:	20979; 20980/25716; 24805
SwissProt:	P21579; Q8N9I0/P46096; P46097/P21707; P29101
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of Synaptotagmin (pT202/199) expression in K562 (A), HEK293T (B), U87MG (C), A549 (D) whole cell lysates.



DEVELOP



IHC description:

Immunohistochemical analysis of Synaptotagmin (pT202/199) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

