

Manufaatuwan/Su	nnligu			
Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82	732223	FAX:	86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name:	Rabbit anti-EMR3 Antibody			
Catalog:	DL94824A			
	22, 102			
Synonyms:	EGF-like module-containing mucin-like hormone			
Synonyms.	receptor-like 3; EGF-like module receptor 3			
	receptor-like 3;	EGF-like m	lodule receptor 3	
		1 .1 .*		
Immunogen:	KLH-conjugated synthetic peptide encompassing a			
	1		region of human EMR3.	
	The exact seque	nce is propi	rietary.	
Form: Liquid		Concentr	ation: 1mg/mL	
Size: 100 ul/50 ul		Host: Rab	bit	
Reactivity: Human, Mouse, Rat		Applicati	on: WB, IF/IC	
Clonality: Polyclonal		Dilution: WB (1/500 - 1/1000),		
• 5				




IF/IC (1/100 - 1/500)

Entrez Gene:	84658
SwissProt:	Q9BY15
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:


Western blot analysis of EMR3 expression in HEK293T (A), H1975 (B), mouse liver (C), mouse kidney (D), rat liver (E), rat kidney (F) whole cell lysates.





IF/ICC description:

Immunofluorescent analysis of EMR3 staining in LOVO cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody i



Storage:



Store at -20°C. Avoid repeated freeze / thaw cycles.

