

Manufacturer/Supplier:			
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD			
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District			
Wuxi Jiangsu Province, China			
TELE: 86-510-82732223 FAX: 86-510-82720101-8014			
Web : <u>www.dldevelop.com</u>			
Email: info@dldevelop.com.cn service@dldevelop.com.cn			
Product name: Catalog:	Rabbit anti-IFNGR1 Antibody DL94847A	r	
Synonyms:	Interferon gamma receptor 1; IFN-gamma receptor 1; IFN-gamma-R1; CDw119; CD119		
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human IFNGR1. The exact sequence is proprietary.		
Form: Liquid	Concentratio	on: 1mg/mL	
Size: 100 ul/50 ul	Host: Rabbit		
Reactivity: Human,Mouse,Rat	<b>Application:</b> WB, IHC, IF/IC at,Monkey		



**Clonality:** Polyclonal

**Dilution:** WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500)


Entrez Gene:	3459/15979
SwissProt:	P15260/P15261
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

#### **WB description:**

Western blot analysis of IFNGR1 expression in Jurkat (A), HepG2 (B), COS7 (C) whole cell lysates.



# **DEVELOP**




#### **IHC description:**

Immunohistochemical analysis of IFNGR1 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

#### **IF/ICC description:**

Immunofluorescent analysis of IFNGR1 staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibod

## **DEVELOP**.



### Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

