

**Manufacturer/Supplier:**

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

**Address:** A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District  
Wuxi Jiangsu Province, China

**TELE:** 86-510-82732223

**FAX:** 86-510-82720101-8014

**Web :** [www.dldevelop.com](http://www.dldevelop.com)

**Email:** [info@dldevelop.com.cn](mailto:info@dldevelop.com.cn)    [service@dldevelop.com.cn](mailto:service@dldevelop.com.cn)

**Product name:** Rabbit anti-Histone H3 Antibody

**Catalog:** DL94866A

**Synonyms:** HIST1H3A; H3FA; HIST1H3B; H3FL;  
HIST1H3C; H3FC; HIST1H3D; H3FB;  
HIST1H3E; H3FD; HIST1H3F; H3FI; HIST1H3G;  
H3FH; HIST1H3H; H3FK; HIST1H3I; H3FF;  
HIST1H3J; H3FJ; Histone H3.1; Histone H3/a;  
Histone H3/b; Histone H3/c; Histone H3/d; Histone  
H3/f; Histone H3/

**Immunogen:** KLH-conjugated synthetic peptide encompassing a  
sequence within the N-term region of human  
Histone H3. The exact sequence is proprietary.

**Form:** Liquid

**Concentration:** 1mg/mL

**Size:** 100 ul/50 ul

**Host:** Rabbit

**Reactivity:**

Human, Mouse, Rat, Bovine, Chicken, Pig, Zebrafish

**Application:** WB, IHC, IF/IC, FC

**Clonality:** Polyclonal

**Dilution:** WB (1/500 - 1/1000),  
IHC (1/100 - 1/200), IF/IC (1/100 -  
1/500), ChIP (Use at an assay  
dependent concentration)

**Entrez Gene:** 319152; 15077; 15078/291159; 100361558

**SwissProt:** P68431; Q71DI3; P84243/P68433; P84228;  
P84244/Q6LED0; P84245


**Purification:** The antibody was purified by immunogen affinity chromatography.

**Buffer:** Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

**WB description:**

Western blot analysis of Histone H3 expression in HEK293T (A), MCF7 (B), Raw264.7 (C), mouse liver (D), rat kidney (E) whole cell

lysates.




### IHC description:

Immunohistochemical analysis of Histone H3 staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

### IF/ICC description:

Immunofluorescent analysis of Histone H3 staining in HEK293T cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary a



**FC description:**

**Storage:**

Store at -20°C. Avoid repeated freeze / thaw cycles.