

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82	2732223	FAX:	86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name:	Rabbit anti-BLK Antibody			
Catalog:	DL95012A			
8				
Synonyms:	Tyrosine-protein kinase Blk; B lymphocyte kinase;			
~JJJ	p55-Blk		-,J	
	pee 2m			
Immunogen:	KLH-conjugated	d synthetic	peptide encompassing a	
g		•		
	sequence within the C-term region of human BLK. The exact sequence is proprietary.			
	The exact seque	nee is prop	netary.	
Form: Liquid		Concentr	ation: 1mg/mL	
Form. Elquid		Concenti	ation. mig/mil	
<b>Size:</b> 100 ul/50 ul		Host: Rab	h;t	
		nost: Kat	oon	
Reactivity: Human		Annligati	on: WB, IHC	
		Applicati	on, wb, mc	
Clonality Dalyala	mal	Dilution	WP(1/500 - 1/1000)	
Clonality: Polyclo	011/01	Dilution:	WB (1/500 - 1/1000),	




## IHC (1/100 - 1/200)

<b>Entrez Gene:</b>	640
SwissProt:	P51451
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

## **WB description:**

Western blot analysis of BLK expression in HepG2 (A), HEK293T (B) whole cell lysates.



## **IHC description:**



Immunohistochemical analysis of BLK staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

## **Storage:**

Store at -20°C. Avoid repeated freeze / thaw cycles.

