

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82732223 FAX: 86-510-82720101-8014			86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: info@dldevelop.com.cn service@dldevelop.com.cn				
	D 111			
Product name:	Rabbit anti-CCR5 Antibody			
Catalog:	DL95249A			
Synonyms:	CMKBR5; C-0	C chemokine	receptor type 5; C-C	
	CKR-5; CC-Cl	KR-5; CCR-	5; CCR5; CHEMR13;	
	HIV-1 fusion c	coreceptor; C	D195	
Immunogen:	KLH-conjugat	ed synthetic	peptide encompassing a	
8	5 0		region of human CCR5.	
	The exact sequ	ence is prop	rietary.	
Form: Liquid		Concentr	ation: 1mg/mL	
Size: 100 ul/50 ul		Host: Rab	bit	
Reactivity: Human, Monkey		Applicati	Application: WB, IHC	



Clonality: Polyclonal

Dilution: WB (1/500 - 1/2000), IHC (1/50 - 1/200)


Entrez Gene:	1234
SwissProt:	P51681
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of CCR5 expression in HEK293T (A) whole cell lysates.



DEVELOP



IHC description:

Immunohistochemical analysis of CCR5 staining in human liver formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.