

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82	2732223	FAX:	86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name:	Rabbit anti-FABP2 Antibody			
Catalog:	DL95250A			
Synonyms:	FABPI; Fatty acid-binding protein, intestinal; Fatty			
	acid-binding pr	otein 2; Inte	stinal-type fatty	
	acid-binding pr	otein; I-FAE	3P	
Immunogen:	KLH-conjugated synthetic peptide encompassing a			
	-		region of human	
	FABP2. The ex	act sequence	e is proprietary.	
		C		
Form: Liquid		Concentr	ation: 1mg/mL	
Size: 100 ul/50 ul		Host: Rab	h;t	
Size: 100 ul/30 ul		Host: Rad	oll	
Reactivity: Human, Mouse		Application	on: WB, IHC	
reactivity. Huma		Philad	···· ·· <i>D</i> , IIIC	



Clonality: Polyclonal

Dilution: WB (1/500 - 1/2000), IHC (1/50 - 1/200)


Entrez Gene:	2169/14079
SwissProt:	P12104/P55050
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of FABP2 expression in K562 (A), mouse lung (B) whole cell lysates.



DEVELOP



IHC description:

Immunohistochemical analysis of FABP2 staining in human small intestine formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.