

Manufacturer/Su	nnlier			
WUXI DONGLIN		VELOPME	NT COLTD	
			ng Road.Liangxi District	
Wuxi Jiangsu Prov	C 1 7			
TELE: 86-510-82	,	FAX:	86-510-82720101-8014	
Web : www.dldev				
Email: info@dldev	•	ervice@dld	levelop.com.cn	
	-	-		
Product name:	Mouse anti-c-FO	OS Antibod	у	
Catalog:	DL95477A			
Synonyms:	G0S7; Proto-one	cogene c-Fo	os; Cellular oncogene	
	fos; G0/G1 swit	ch regulator	ry protein 7	
Immunogen:	-	otein corres	sponding to human	
	c-FOS.			
Form: Liquid		Concentr	ation: 1mg/mL	
C* 100 1/50 1				
Size: 100 ul/50 ul		Host: Mo	use	
Reactivity: Huma	n Mouso Pot	Applicati	on: WB, IHC	
Reactivity. IIuilla	II,Iviouse,Rat	Applicati	on. wb, mc	
Clonality: Monoc	Clonality: Monoclonal Dilution: WB (1/1000 - 1/2000),			
Cronunty, monou	ionul	IHC (1/50		
		(1.00	=,	

DEVELOP

Entrez Gene: 23


SwissProt: P01100

Purification:

Buffer: Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of c-FOS expression in Hela (A), 293T (B), mouse brain (C), rat brain (D) whole cell lysates.



IHC description:

Immunohistochemical analysis of c-FOS staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was



pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

