

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

Web : www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Mouse anti-NF-kappaB p65 Antibody

Catalog: DL95538A

Synonyms: NFKB3; Transcription factor p65; Nuclear factor
NF-kappa-B p65 subunit; Nuclear factor of kappa
light polypeptide gene enhancer in B-cells 3

Immunogen: KLH-conjugated synthetic peptide encompassing a
sequence of human NF-kappaB p65. The exact
sequence is proprietary.

Form: Liquid

Concentration: 1mg/mL

Size: 100 ul/50 ul

Host: Mouse

Reactivity: Human, Mouse, Rat

Application: WB, IHC, IP

Clonality: Monoclonal

Dilution: WB (1/1000 - 1/3000),
IHC (1/100 - 1/200), IP (1/50 -
1/200)

Entrez Gene: 5970/19697


SwissProt: Q04206/Q04207

Purification:

Buffer: Liquid in 0.42% Potassium phosphate, 0.87%
Sodium chloride, pH 7.3, 30% glycerol, and 0.01%
sodium azide.

WB description:

Western blot analysis of NF-kappaB p65 expression in Hela (A),
293T (B), HepG (C) whole cell lysates.




IHC description:

Immunohistochemical analysis of NF-kappaB p65 staining in mouse hippocampus formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IP description:

Immunoprecipitation of NF-kappaB p65 from 0.5mg Hela whole cell extract lysate, using anti-NF-kappaB p65 Antibody.



Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.