

Manufacturer/Supplier:			
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD			
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District			
Wuxi Jiangsu Province, China			
TELE: 86-510-82732223			86-510-82720101-8014
Web : <u>www.dldevelop.com</u>			
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>			
Product name:	Mouse anti-OCT1 Antibody		
Catalog:	DL95539A		
Synonyms:	OCT1; OTF1; POU domain class 2 transcription		
	factor 1; NF-A1; Octamer-binding protein 1; Oct-1;		
	Octamer-binding transcription factor 1; OTF-1		
Immunogen:	KLH-conjugated synthetic peptide encompassing a		
	sequence of human OCT1. The exact sequence is		
	proprietary.		
T. T. 1		C (	· 1 / T
Form: Liquid		Concentra	ation: 1mg/mL
Size: 100 u1/50 u1		Host: Mo	
Size. 100 ul/30 ul		HOSE: MO	use
Reactivity: Human		Application: WB, IHC	
reactivity. Human		rppncau	, , , , , , , , , , , , , , , , , , ,



**Clonality:** Monoclonal

**Dilution:** WB (1/500 - 1/2000), IHC (1/50 - 1/200)

**Entrez Gene:** 


**SwissProt:** P14859; P09086

**Purification:** 

Buffer: Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

## WB description:

Western blot analysis of 42278 expression in Hela (A), Jurkat (B), HepG2 (C) whole cell lysates.



**IHC description:** 



Immunohistochemical analysis of 42278 staining in human tonsil formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

## **Storage:**

Store at -20°C. Avoid repeated freeze / thaw cycles.

