

Manufacturer/Supplier:			
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD			
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District			
Wuxi Jiangsu Province, China			
TELE: 86-510-82732223 FAX: 86-510-82720101-8014			
Web : <u>www.dldevelop.com</u>			
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>			
Product name:	Rabbit anti-CD184 Antibody		
Catalog:	DL95604A		
Synonyms:	C-X-C chemokine receptor type 4; CXC-R4;		
	CXCR-4; FB22; Fusin; HM89; LCR1; Leukocyte-derived seven transmembrane domain receptor; LESTR; Lipopolysaccharide-associated		
	protein 3; LAP-3; LPS-associated protein 3;		
	NPYRL; Stromal cell-derived factor 1 receptor		
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence of human CD184. The exact sequence is		
	proprietary.		
Form: Liquid	Concentration: 1mg/mL		
Size: 100 ul/50 ul	Host: Rabbit		




Reactivity: Human, Mouse, Rat		Application: WB, IHC
Clonality: Polyclonal		<b>Dilution:</b> WB (1/1000 - 1/2000), IHC (1/100 - 1/200)
Entrez Gene:	7852	
SwissProt:	P61073	
Purification:	The antibody was purified by immunogen affinity chromatography.	
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.	

## **WB description:**

Western blot analysis of CD184 expression in Hela (A), rat brain (B) whole cell lysates.

## **DEVELOP**



## **IHC description:**

Immunohistochemical analysis of CD184 staining in human breast cancer,human tonsil formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

## **Storage:**

Store at -20°C. Avoid repeated freeze / thaw cycles.