

Manufacturer/Supplier:			
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD			
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District			
Wuxi Jiangsu Province, China			
TELE: 86-510-82	FAX: 86-510-82720101-8014		
Web : <u>www.dldevelop.com</u>			
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>			
Product name:	Rabbit anti-Histone H3 (TriMethyl K18) Antibody		
Catalog:	DL95712A		
Synonyms:	HIST1H3A; H3FA; HIST1H3B; H3FL; HIST1H3C; H3FC; HIST1H3D; H3FB;		
	HIST1H3E; H3FD; HIST1H3F; H3FI; HIST1H3G;		
	H3FH; HIST1H3H; H3FK; HIST1H3I; H3FF;		
	HIST1H3J; H3FJ; Histone H3.1; Histone H3/a;		
	Histone H3/b; Histone H3/c; Histone H3/d; Histone		
	H3/f; Histone H3/		
Immunogen:	KLH-conjugated synthetic peptide encompassing a		
	sequence of human Histone H3. The exact sequence		
	is proprietary.		
Form: Liquid	Concentration: lmg/mL		




Size: 100 ul/50 ul		Host: Rabbit
Reactivity: Human, Mouse		Application: WB, IHC
Clonality: Polyclonal		Dilution: WB (1/500 - 1/1000), IHC (1/200 - 1/500)
Entrez Gene:	319152; 15077;	15078
SwissProt:	P68431; Q71DI3; P84243/P68433; P84228; P84244	
Purification:	The antibody was purified by immunogen affinity chromatography.	
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.	

WB description:

Western blot analysis of Histone H3 (TriMethyl K18) expression in Hela (A), NIH3T3 (B) whole cell lysates.

DEVELOP



IHC description:

Immunohistochemical analysis of Histone H3 (TriMethyl K18) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

