

Manufacturer/Supplier:			
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD			
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District			
Wuxi Jiangsu Province, China			
TELE: 86-510-82	FAX: 86-510-82720101-8014		
Web : <u>www.dldevelop.com</u>			
Email: info@dldevelop.com.cn service@dldevelop.com.cn			
Product name:	Rabbit anti-LC3B Antibody		
Catalog:	DL95753A		
Synonyms:	MAP1ALC3; Microtubule-associated proteins 1A/1B light chain 3B; Autophagy-related protein LC3 B; Autophagy-related ubiquitin-like modifier		
	LC3 B; MAP1 light chain 3-like protein 2;		
	MAP1A/MAP1B light chain 3 B; MAP1A/MAP1B		
	LC3 B; Microtubule-associated pro		
Immunogen:	Recombinant protein corresponding to human		
	LC3B.		
Form: Liquid	Concentration: 1mg/mL		
Size: 100 ul/50 ul	Host: Rabbit		




Reactivity: Human, Mouse, Rat		Application: WB, IHC, IF/IC
Clonality: Polyclo	nal	<b>Dilution:</b> WB (1/1000 - 1/2000), IHC (1/200 - 1/500), IF/IC (1/100 - 1/200)
Entrez Gene:	81631	
SwissProt:	Q9GZQ8	
Purification:	The antibody was purified by immunogen affinity chromatography.	
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.	

#### **WB description:**

Western blot analysis of LC3B expression in Hela (A), NIH3T3 (B), rat brain (C) whole cell lysates.



# **DEVELOP**




#### **IHC description:**

Immunohistochemical analysis of LC3B staining in rat brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

### **IF/ICC description:**

Immunofluorescent analysis of LC3B staining in Hela cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody i

## **DEVELOP**.



#### Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

