

## Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

**Web**: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Rabbit anti-MKK1 Antibody

**Catalog:** DL95758A

MEK1; PRKMK1; Dual specificity **Synonyms:** 

> mitogen-activated protein kinase kinase 1; MAP kinase kinase 1; MAPKK 1; MKK1; ERK activator

kinase 1; MAPK/ERK kinase 1; MEK 1

Immunogen: Recombinant protein corresponding to human

MKK1.

Form: Liquid Concentration: 1mg/mL

**Size:** 100 ul/50 ul Host: Rabbit

Reactivity: Human, Mouse, Rat **Application:** WB, IHC



**Clonality:** Polyclonal **Dilution:** WB (1/1000 - 1/2000),

IHC (1/100 - 1/200)

**Entrez Gene:** 5604

Q02750 SwissProt:

**Purification:** The antibody was purified by immunogen affinity

chromatography.

**Buffer:** Liquid in 0.42% Potassium phosphate, 0.87%


Sodium chloride, pH 7.3, 30% glycerol, and 0.01%

sodium azide.

**WB** description:

Western blot analysis of MKK1 expression in Hela (A), NIH3T3 (B), PC12 (C) whole cell lysates.





## **IHC** description:

Immunohistochemical analysis of MKK1 staining in human colon cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

## **Storage:**

Store at -20°C. Avoid repeated freeze / thaw cycles.