

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTDAddress: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi DistrictWuxi Jiangsu Province, ChinaTELE: 86-510-82732223FAX: 86-510-82720101-8014Web : www.dldevelop.comEmail: info@dldevelop.com.cnservice@dldevelop.com.cn		
Wuxi Jiangsu Province, China TELE: 86-510-82732223 FAX: 86-510-82720101-8014 Web: www.dldevelop.com		
TELE: 86-510-82732223 FAX: 86-510-82720101-8014 Web : www.dldevelop.com		
Web : <u>www.dldevelop.com</u>		
Email: info@dldevelop.com.cn service@dldevelop.com.cn		
Product name: Rabbit anti-TBP Antibody		
Catalog: DL95794A		
Synonyms: GTF2D1; TF2D; TFIID; TATA-box-binding		
protein; TATA sequence-binding protein;		
TATA-binding factor; TATA-box factor;		
Transcription initiation factor TFIID TBP subunit		
Immunogen: Recombinant protein corresponding to human TBP.		
Form: Liquid Concentration: 1mg/mL		
Concentration. Ting inc		
Size: 100 ul/50 ul Host: Rabbit		
Reactivity: Human, Mouse, Rat Application: WB, IHC		
Clonality: Polyclonal Dilution: WB (1/1000 - 1/2000),		




IHC (1/200 - 1/500)

Entrez Gene:	6908
SwissProt:	P20226
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of TBP expression in Hela (A), 293T (B), C2C12 (C), mouse liver (D), PC12 (E), rat brain (F) whole cell lysates.



IHC description:



Immunohistochemical analysis of TBP staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

