

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Rabbit anti-DCK Antibody

Catalog: DL95942A

Synonyms: Deoxycytidine kinase; dCK

Immunogen: Recombinant full length protein of human DCK

Form: Liquid Concentration: 1mg/mL

Size: 100 ul/50 ul Host: Rabbit

Reactivity: Human, Mouse, Rat **Application:** WB, IHC, IF/IC

Clonality: Polyclonal Dilution: WB (1/500 - 1/2000),

IHC (1/50 - 1/200), IF/IC (1/50 -

1/200)



Entrez Gene: 1633/13178/79127

P27707/P43346/P48769 SwissProt:

Purification: The antibody was purified by immunogen affinity

chromatography.


Buffer: Liquid in 0.42% Potassium phosphate, 0.87%

Sodium chloride, pH 7.3, 30% glycerol, and 0.01%

sodium azide.

WB description:

Western blot analysis of DCK expression in Raji (A), HepG2 (B) whole cell lysates.



IHC description:


Immunohistochemical analysis of DCK staining in rat kidney formalin fixed paraffin embedded tissue section. The section was



pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IF/ICC description:

Immunofluorescent analysis of DCK staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in



Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.