

**Manufacturer/Supplier:**

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

**Address:** A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District  
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

**Web :** [www.dldevelop.com](http://www.dldevelop.com)

Email: [info@dldevelop.com.cn](mailto:info@dldevelop.com.cn)    [service@dldevelop.com.cn](mailto:service@dldevelop.com.cn)

**Product name:** Rabbit anti-Reptin 52 Antibody

**Catalog:** DL95953A

**Synonyms:** INO80J; TIP48; TIP49B; RuvB-like 2; 48 kDa  
TATA box-binding protein-interacting protein; 48  
kDa TBP-interacting protein; 51 kDa erythrocyte  
cytosolic protein; ECP-51; INO80 complex subunit  
J; Repressing pontin 52; Reptin 52; TIP49b;  
TIP60-associated prote

**Immunogen:** Recombinant full length protein of human Reptin 52

**Form:** Liquid

**Concentration:** 1mg/mL

**Size:** 100 ul/50 ul

**Host:** Rabbit

**Reactivity:** Human, Mouse, Rat

**Application:** WB, IF/IC

**Clonality:** Polyclonal

**Dilution:** WB (1/500 - 1/2000),  
IF/IC (1/50 - 1/200)

**Entrez Gene:** 10856/20174


**SwissProt:** Q9Y230/Q9WTM5

**Purification:** The antibody was purified by immunogen affinity chromatography.

**Buffer:** Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.


**WB description:**

Western blot analysis of Reptin 52 expression in K562 (A), SW480 (B), MCF7 (C) whole cell lysates.



### IF/ICC description:

Immunofluorescent analysis of Reptin 52 staining in U2OS cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody.



### Storage:



Store at -20°C. Avoid repeated freeze / thaw cycles.