


Manufacturer/Supplier: WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82732223		FAX:	86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name:	Rabbit anti-Histone H2B Antibody			
Catalog:	DL95962A			
Synonyms:	HODEO, Histons HOD type 2 E. Histons			
Synonyms.	H2BFQ; Histone H2B type 2-E; Histone			
	H2B-GL105; Hi	Istone H2B	q; H2B/q	
Immunogen:	Recombinant full length protein of human Histone			
	H2B			
Form: Liquid		Concentr	ation: 1mg/mL	
Size: 100 ul/50 ul		Host: Rabbit		
Reactivity: Human, Mouse, Rat Application: W			we WB IHC	
Reactivity. Human, wouse, Rat		Аррисан	on, wb, me	
Clansity Debuland Dilution WD (1/500 1/2000)				
Clonality: Polyclonal		Dilution: WB (1/500 - 1/2000),		
		IHC (1/50	- 1/200)	

DEVELOP.

Entrez Gene:	8349/319190
SwissProt:	Q16778/Q64524
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of Histone H2B expression in SW480 (A), BT474 (B) whole cell lysates.



IHC description:

Immunohistochemical analysis of Histone H2B staining in human



esophageal cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

