

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Rabbit anti-Eppin Antibody

Catalog: DL95986A

Synonyms: SPINLW1; WAP7; WFDC7; Eppin; Cancer/testis

antigen 71; CT71; Epididymal protease inhibitor;

Protease inhibitor WAP7; Serine protease

inhibitor-like with Kunitz and WAP domains 1:

WAP four-disulfide core domain protein 7

Recombinant full length protein of human Eppin Immunogen:

Form: Liquid Concentration: 1mg/mL

Size: 100 u1/50 u1 Host: Rabbit

Reactivity: Human, Mouse **Application:** WB, IF/IC



Clonality: Polyclonal **Dilution:** WB (1/500 - 1/2000),

IF/IC (1/50 - 1/200)

Entrez Gene: 75526

O95925/O9DA01 SwissProt:

Purification: The antibody was purified by immunogen affinity

chromatography.

Buffer: Liquid in 0.42% Potassium phosphate, 0.87%


Sodium chloride, pH 7.3, 30% glycerol, and 0.01%

sodium azide.

WB description:


Western blot analysis of Eppin expression in THP1 (A), mouse liver (B), mouse testis (C) whole cell lysates.





IF/ICC description:

Immunofluorescent analysis of Eppin staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody



Storage:



Store at -20°C. Avoid repeated freeze / thaw cycles.