

## Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>

**Product name:** Rabbit anti-DDB1 Antibody

Catalog: DL96015A

**Synonyms:** XAP1; DNA damage-binding protein 1; DDB p127

subunit; DNA damage-binding protein a; DDBa;
Damage-specific DNA-binding protein 1; HBV
X-associated protein 1; XAP-1; UV-damaged
DNA-binding factor; UV-damaged DNA-binding

protein 1; UV-DDB 1; XPE-binding fact

Immunogen: KLH-conjugated synthetic peptide of human DDB1

Form: Liquid Concentration: 1mg/mL

Size: 100 ul/50 ul Host: Rabbit

Reactivity: Human, Mouse, Rat Application: WB, IHC



**Clonality:** Polyclonal **Dilution:** WB (1/500 - 1/1000),

IHC (1/50 - 1/100)

Entrez Gene: 1642/13194

Q16531/Q3U1J4/Q9ESW0 SwissProt:

**Purification:** The antibody was purified by immunogen affinity

chromatography.

**Buffer:** Liquid in 0.42% Potassium phosphate, 0.87%


Sodium chloride, pH 7.3, 30% glycerol, and 0.01%

sodium azide.

## **WB** description:

Western blot analysis of DDB1 expression in A549 (A), mouse brain (B), mouse spleen (C), mouse liver (D) whole cell lysates.





## **IHC** description:

Immunohistochemical analysis of DDB1 staining in human kidney formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

## **Storage:**

Store at -20°C. Avoid repeated freeze / thaw cycles.