


Manufaaturar/Su					
Manufacturer/Supplier: WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD					
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District					
Wuxi Jiangsu Province, China					
TELE: 86-510-82732223		FAX:	86-510-82720101-8014		
Web : <u>www.dldevelop.com</u>					
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>					
Product name:	Rabbit anti-Rabphilin-3AL Antibody				
Catalog:	DL96036A				
8					
Synonyms:	NOC2; Rab effector Noc2; No C2 domains protein;				
Synonyms.	Rabphilin-3A-li		rto c2 domanis protein,		
	Kaopinini-5A-n	ke protein			
T	D	11 1 41			
Immunogen:	Recombinant full length protein of human				
	Rabphilin-3AL				
Form: Liquid		Concentr	ation: 1mg/mL		
Size: 100 ul/50 ul		Host: Rab	bit		
Reactivity: Human		Applicati	on: WB, IHC		
Clonality: Polyclonal		Dilution:	WB (1/500 - 1/2000),		
		IHC (1/50			
		(0 0	/		

www.dldevelop.com

Entrez Gene:	9501
SwissProt:	Q9UNE2
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of Rabphilin-3AL expression in BT474 (A), 22RV1 (B), SW620 (C) whole cell lysates.



IHC description:

Immunohistochemical analysis of Rabphilin-3AL staining in

www.dldevelop.com



human stomach formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.