

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Rabbit anti-UQCR10 Antibody

Catalog: DL96038A

Synonyms: UCRC; Cytochrome b-c1 complex subunit 9;

Complex III subunit 9; Complex III subunit X;

Cytochrome c1 non-heme 7 kDa protein;

Ubiquinol-cytochrome c reductase complex 7.2 kDa

protein

Recombinant full length protein of human UQCR10 Immunogen:

Form: Liquid Concentration: 1mg/mL

Size: 100 u1/50 u1 Host: Rabbit

Reactivity: Human **Application:** WB, IHC



Clonality: Polyclonal **Dilution:** WB (1/500 - 1/2000),

IHC (1/50 - 1/200)

Entrez Gene: 29796

O9UDW1 SwissProt:

Purification: The antibody was purified by immunogen affinity

chromatography.

Buffer: Liquid in 0.42% Potassium phosphate, 0.87%


Sodium chloride, pH 7.3, 30% glycerol, and 0.01%

sodium azide.

WB description:

Western blot analysis of UQCR10 expression in MCF7 (A), Jurkat (B) whole cell lysates.





IHC description:

Immunohistochemical analysis of UQCR10 staining in human liver cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.