

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82732223 FAX: 86-510-82720101-8014				
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name:	Rabbit anti-Interferon lambda-3 Antibody			
Catalog:	DL96109A			
Synonyms:	IL28B; IL28C; ZCYTO22; Interferon lambda-3;			
		•	cyto22; Interleukin-28B;	
	IL-28B; Interleu	ıkin-28C; Il	L-28C	
T	Decembra de Co	11 1 41		
Immunogen:	Recombinant full length protein of human Interferon lambda-3			
	Interferon lambo	ua-3		
Form: Liquid		Concentr	ation: 1mg/mL	
I of m. Erquite		concenti		
Size: 100 ul/50 ul		Host: Rabbit		
Reactivity: Human		Application: WB, IHC		
Clonality: Polyclonal		Dilution: WB (1/500 - 1/2000),		




IHC (1/50 - 1/200)

Entrez Gene:	282617
SwissProt:	Q8IZI9
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of Interferon lambda-3 expression in HEK293T (A) whole cell lysates.



IHC description:



Immunohistochemical analysis of Interferon lambda-3 staining in rat brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

