


Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82732223		FAX:	86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: info@dldevelop.com.cn service@dldevelop.com.cn				
	-	-	-	
Product name:	Rabbit anti-SNRPA Antibody			
Catalog:	DL96232A			
Catalog.	DL)0232A			
Synonyms:		r ribonucle	oprotein A; U1 snRNP	
	A; U1-A; U1A			
Immunogen:	Recombinant full length protein of human SNRPA			
Form: Liquid		Concentra	ation: 1mg/mL	
Size: 100 ul/50 ul		Host: Rabbit		
		110000 1100		
Reactivity: Human, Mouse, Rat Application: WB, IHC, IF/IC				
Reactivity: Human, Mouse, Rat		Application	<b>III.</b> WD, IIIC, II/IC	
(1, 1) $(1, 2)$ $(1, 2)$				
<b>Clonality:</b> Polyclonal		<b>Dilution:</b> WB (1/500 - 1/2000),		
			- 1/200), IF/IC (1/50 -	
		1/200)		

# **DEVELOP**

Entrez Gene:	6626/53607
SwissProt:	P09012/Q62189
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

## **WB description:**

Western blot analysis of SNRPA expression in A431 (A), MCF7 (B), mouse testis (C), rat brain (D) whole cell lysates.



# **IHC description:**


Immunohistochemical analysis of SNRPA staining in mouse brain



formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

### **IF/ICC description:**

Immunofluorescent analysis of SNRPA staining in Hela cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody



#### **Storage:**

Store at -20°C. Avoid repeated freeze / thaw cycles.