


Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82	2732223	FAX:	86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name: Catalog:	Rabbit anti-eIF2 DL96277A	2S3 Antiboo	ły	
Synonyms:	EIF2G; Eukaryotic translation initiation factor 2 subunit 3; Eukaryotic translation initiation factor 2 subunit gamma X; eIF-2-gamma X; eIF-2gX			
Immunogen:	Recombinant full length protein of human eIF2S3			
Form: Liquid		Concentr	ation: 1mg/mL	
Size: 100 ul/50 ul		Host: Rat	bit	
Reactivity: Human, Mouse, Rat		Application: WB, IHC		
Clonality: Polyclonal		Dilution: WB (1/500 - 1/2000), IHC (1/50 - 1/200)		



Entrez Gene:	1968/26905/100312984; 299027
SwissProt:	P41091/Q9Z0N1/P81795
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of eIF2S3 expression in SW620 (A), HL60 (B) whole cell lysates.



IHC description:

Immunohistochemical analysis of eIF2S3 staining in rat brain

www.dldevelop.com



formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.